

9/10

## Section 12.5: Lines & Planes

A line in 3-space is given by parametrized vector equation:  $\boxed{l(t) = \vec{p} + t\vec{v}}$  where  $\vec{p}$  = position vector of a point on  $l$  and  $\vec{v}$  = direction of line

Ex: Compute the vector equation of the line through  $(-6, 2, 3)$  and parallel to line  $m(t) = \langle 0, 2, -1 \rangle + t\langle -2, 1, 5 \rangle$

Sol Given:  $\vec{p} = \langle -6, 2, 3 \rangle$ . Because of parallelism,  $\vec{v} = \langle -2, 1, 5 \rangle$  is a valid direction vector.

$$\therefore l(t) = \langle -6, 2, 3 \rangle + t\langle -2, 1, 5 \rangle$$

The parametric equations of a line are  $\begin{cases} x = x(t) \\ y = y(t) \\ z = z(t) \end{cases}$  which are the component functions of the vector form

Ex: For  $l$  and  $m$  as in the previous example, simplify the vector equation:

$$l(t) = (-6, -2t, 2+t, 3+5t) \quad \begin{cases} x = -6-2t \\ y = 2+t \\ z = 3+5t \end{cases}$$

$$m(t) = (-2t, 2+t, -1+5t) \quad \begin{cases} x = -2t \\ y = 2+t \\ z = -1+5t \end{cases}$$

$\therefore$  has parametric equations

A line can also be represented (sometimes) by symmetric equations

(solved for parameter)

$$\frac{x-x_0}{a} = \frac{y-y_0}{b} = \frac{z-z_0}{c}$$

Ex: For  $l$  as above, has parametric equations:

$$\begin{cases} x = -6-2t \\ y = 2+t \\ z = 3+5t \end{cases}$$

$$\Rightarrow \begin{cases} \frac{x+6}{-2} = t \\ \frac{y-2}{1} = t \\ \frac{z-3}{5} = t \end{cases}$$

$$\frac{x+6}{-2} = \frac{y-2}{1} = \frac{z-3}{5}$$

Symmetric equations of  $l$

This procedure can be done in reverse too; if given symmetry equations we can easily get parametric equations too

Some Terminology: Two lines are ...

- ① parallel if their direction vectors are parallel
- ② intersecting if they have a point in common
- ③ skew if they are neither parallel or intersecting

Ex: Classify as parallel, intersecting, or skew:

$$l_1(t) = \langle 5 - 12t, 3 + 9t, 1 - 3t \rangle$$

$$l_2(t) = \langle 3 + 8t, -6t, 7 + 2t \rangle$$

sol: 
$$\left. \begin{aligned} l_1(t) &= \langle 5, 3, 1 \rangle + t \langle -12, 9, -3 \rangle \\ l_2(t) &= \langle 3, 0, 7 \rangle + t \langle 8, -6, 2 \rangle \end{aligned} \right\} \text{not equal, but we must check the unit vectors}$$

unit vectors:

$$\frac{1}{|\vec{v}_1|} \vec{v}_1 = \frac{1}{\sqrt{12+9+9}} \langle -12, 9, -3 \rangle = \frac{1}{3\sqrt{26}} \langle -12, 9, -3 \rangle = \boxed{\frac{1}{\sqrt{26}} \langle -4, 3, -1 \rangle}$$

$$\frac{1}{|\vec{v}_2|} \vec{v}_2 = \frac{1}{\sqrt{64+36+4}} \langle 8, -6, 2 \rangle = \frac{1}{2\sqrt{26}} \langle 8, -6, 2 \rangle = \boxed{\frac{1}{\sqrt{26}} \langle 4, -3, 1 \rangle}$$

Notice:  $\frac{1}{|\vec{v}_2|} \vec{v}_2 = -\frac{1}{|\vec{v}_1|} \vec{v}_1$ , so  $l_1$  is parallel to  $l_2$

check if they're intersecting:

Do NOT solve  $l_1(t) = l_2(t)$ . Instead, solve for something like  $l_1(t) = l_2(s) \Rightarrow$  do paths cross, not necessarily at the same time:

$$\langle 5 - 12t, 3 + 9t, 1 - 3t \rangle = \langle 3 + 8s, -6s, 7 + 2s \rangle$$

ie. 
$$\begin{cases} 5 - 12t = 3 + 8s \\ 3 + 9t = -6s \\ 1 - 3t = 7 + 2s \end{cases} \Rightarrow \begin{cases} -12t - 8s = -2 \\ 9t + 6s = -3 \\ -3t - 2s = 6 \end{cases} \Rightarrow \text{This is a system of equations}$$



Ex Cont:  $\begin{cases} 6t+4s=1 \\ 3t+2s=-1 \\ 3t+2s=-6 \end{cases} \Rightarrow$  implies  $-1 = 3t+2s = -6$  which is simply not a valid expression.

$\therefore$  the lines are not intersecting

Recall: A plane in 3-space has vector equation

$$\vec{n} \cdot (\vec{x} - \vec{p}) = 0$$

normal vector  $\rightarrow \vec{n}$        $\vec{x}$  vector of variables       $\vec{p}$  any position point on the plane

Ex: Compute the plane through  $(1, 2, 4)$  and perpendicular to  $\langle -2, 1, 3 \rangle$ :


$$\vec{n} \cdot (\vec{x} - \vec{p}) = 0 \Rightarrow \langle -2, 1, 3 \rangle \cdot \langle x-1, y-2, z-4 \rangle = 0$$

$$= -2(x-1) + 1(y-2) + 3(z-4) = 0$$

Ex: Compute the plane through the point  $(3, 5, -1)$  and containing the line  $\begin{cases} x=4-t \\ y=2t-1 \\ z=-3t \end{cases}$

Sol:  $\vec{p} = (3, 5, -1)$

need  $Q$ , a point on  $l \Rightarrow$  pick a time  
so let's use  $Q = l(0) = (4, -1, 0)$



So,  $\vec{n} = (3-4, 5-(-1), -1-0) = \langle -1, 6, -1 \rangle$

$$l(t) = (4, -1, 0) + \underbrace{\langle -1, 0, -3 \rangle}_{\vec{v}}$$

Ex Cont:  $\vec{n} = \vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -1 & 6 & -1 \\ -1 & 2 & -3 \end{vmatrix} = \langle -16, -2, 4 \rangle$

$\therefore$  the plane has equation  $\vec{n} \cdot (\vec{x} - \vec{p}) = 0$

i.e.  $\langle -16, -2, 4 \rangle \cdot \langle x-3, y-5, z+1 \rangle = 0$   
 $\Rightarrow -16(x-3) - 2(y-5) + 4(z+1) = 0$

## Section 12.6: Quadratic Surfaces

IDEA: We want to study degree 2 polynomials and solution sets in 3-space

Ex:  $p(x, y, z) = x^2 - z \leftarrow$  "degenerate" b/c it doesn't depend on all variables

solution set:  $p(x, y, z) = 0$  iff  $x^2 - z = 0 \Rightarrow x^2 = z$

In the  $xz$  plane, this looks like a parabola

This solution set is actually a (parabolic) cylinder

A picture in 3-space:

